



**education**

Department:  
Education  
PROVINCE OF KWAZULU-NATAL

**NATIONAL  
SENIOR CERTIFICATE**

**GRADE 11**

**PHYSICAL SCIENCES P2  
(CHEMISTRY)**

**COMMON TEST**

**JUNE 2019**

**MARKS: 100**

**TIME : 2 Hours**

**This question paper consists of 9 pages and 2 data sheets.**

**INSTRUCTIONS AND INFORMATION TO CANDIDATES**

1. Write your name on the **ANSWER BOOK**.
2. Answer **ALL** the questions in the answer book.
3. You may use a non-programmable calculator.
4. You may use appropriate mathematical instruments.
5. Number the answers correctly according to the numbering system used in this question paper.
6. **YOU ARE ADVISED TO USE THE ATTACHED DATA SHEETS.**
7. Give brief motivations, discussions, et cetera where required.
8. Show the formulae and substitutions in **ALL** calculations.
9. Round off answers to a minimum of **TWO** decimal places

**QUESTION 1: MULTIPLE- CHOICE QUESTIONS**

Four options are provided as possible answers to the following questions. Each question has only ONE correct answer. Write only the letter (A - D) next to the question number (1.1 – 1.7) in the ANSWER BOOK. Eg 1.8 A

1.1 The shape of the  $\text{OF}_2$  molecule is :

- A Trigonal pyramidal
- B Linear
- C Bent
- D Tetrahedral

(2)

1.2 Which statement BEST explains the formation of the dative bond between ammonia ( $\text{NH}_3$ ) and the hydrogen ion ( $\text{H}^+$ )?

- A Both  $\text{NH}_3$  and  $\text{H}^+$  are polar
- B The  $\text{NH}_3$  molecule has a lone pair of electrons and the  $\text{H}^+$  ion has an empty orbital.
- C  $\text{H}^+$  ion is regarded as a proton and is attracted to the electrons on the nitrogen atom of the  $\text{NH}_3$  molecule.
- D The electronegativity of the nitrogen atom is greater than the electronegativity of hydrogen.

(2)

1.3 Which ONE of the following statements concerning ideal gases is **INCORRECT**?

- A Ideal gases do not exert pressure
- B Ideal gas molecules do not occupy a volume
- C The collision between ideal gas molecules is elastic
- D There are no intermolecular forces between ideal gas molecules.


(2)

1.4 Which ONE of the following statements regarding the effect of intermolecular forces and some physical properties is **INCORRECT**?

- A The stronger the intermolecular force, the slower the rate of evaporation.
- B The weaker the intermolecular force, the lower the boiling point.
- C The stronger the intermolecular force, the higher the surface tension.
- D The stronger the intermolecular force, the lower the melting point..

(2)

- 1.5 2,50 mol of  $\text{SO}_2$  and 1 mol of  $\text{O}_2$  are sealed in a  $1 \text{ dm}^3$  flask and allowed to react completely at STP according to the following balanced equation.



The TOTAL number of moles of gas in the flask at the END of the reaction is:

- A 2  
B 3,50  
C 2,50  
D 0,50 (2)
- 1.6 The gas law that expresses the relationship between pressure and temperature of a gas is known as:
- A Charle's Law  
B Gay- Lussac Law  
C Boyle's Law  
D Avogadro's Law (2)
- 1.7 The type of intermolecular force involved when  $\text{CO}_2$  is added to water is :
- A Dipole- induced dipole forces  
B London forces  
C Covalent bonds  
D Ion – induced dipole forces (2)

**TOTAL : SECTION A [14]**

**QUESTION 2**

Consider the substances in the table below. Select the correct answer for each of the questions that follow. Write down only the LETTER that corresponds to your choice

	SUBSTANCE		SUBSTANCE
A.	HCN	F	H <sub>2</sub> O
B	MgCl <sub>2</sub>	G	CCl <sub>4</sub>
C	I <sub>2</sub>	H	CO <sub>2</sub>
D	NH <sub>4</sub> <sup>+</sup>	I	C <sub>2</sub> H <sub>2</sub>
E	Cl <sub>2</sub>	J	H <sub>2</sub> S

**2.1 Identify**

- 2.1.1 TWO molecules that have triple bonds. (2)
- 2.1.2 TWO substances that when mixed together will result in ion-dipole forces of attraction. (2)
- 2.1.3 A non-polar LIQUID at room temperature (1)
- 2.1.4 A MOLECULE having a tetrahedral shape. (1)
- 2.1.5 A SOLID that is insoluble in water. (1)
- 2.1.6 A GAS at room temperature with pure covalent bonds between its atoms. (1)

- 2.2 Compounds F (H<sub>2</sub>O) and J (H<sub>2</sub>S) are hydrides of group 6 elements. H<sub>2</sub>O has a lower molar mass than H<sub>2</sub>S, but a higher boiling point than H<sub>2</sub>S.

Explain fully why H<sub>2</sub>O has a higher boiling point than H<sub>2</sub>S, by referring to the types and strengths of the intermolecular forces in each and the energy involved. (4)

- 2.3 Draw the Lewis structure for compound G (CH<sub>4</sub>) (2)

- 2.4 Compound H (CO<sub>2</sub>) has polar covalent bonds in the molecule.


2.4.1 Is the compound CO<sub>2</sub> polar or non-polar? Explain fully. (3)

2.4.2 Name the type of intermolecular forces found in this compound. (1)

[18]

**QUESTION 3**

Grade 11 learners investigated the effect of intermolecular force on capillarity. They pour 100ml each of **water; glycerine and nail polish remover** in separate beakers. A capillary tube is inserted into each liquid and after a while, the level of liquid in the capillary tube is measured.



They recorded their results in a table as follows:


Liquid	Height (mm)
Water	19
Glycerine	5
Nail polish remover	26

- 3.1 State the dependent variable in the above investigation. (1)
- 3.2 Which liquid displayed the greatest degree of capillarity? (1)
- 3.3 Explain the answer to question 3.2 above. (3)
- 3.4 Use the results in the table and arrange the liquids in order of **INCREASING STRENGTH** of intermolecular force that is, from the weakest intermolecular force to the strongest intermolecular force. (2)
- 3.5 Identify the liquid with the highest boiling point. (1)
- [8]**

**QUESTION 4**

- 4.1 The graph below shows the relationship between the volume and Celsius temperature of an enclosed gas maintained at a constant pressure,  $P_1$ . The experiment is repeated for different constant pressures  $P_2$  and  $P_3$ .

**Graph of volume against temperature**



- 4.1.1 Give the name, and state in words, the Law that is illustrated in the graph. (3)
- 4.1.2 Apart from pressure, state ONE other variable that must be kept constant for each experiment (1)
- 4.1.3 What is the relationship between the volume and temperature of the gas? (1)
- 4.1.4 Which one of the 3 pressures,  $P_1$ ;  $P_2$  or  $P_3$  is the HIGHEST ? (1)
- 4.1.5 Fully explain the answer to question 4.1.4 above. Use a relevant equation to support your explanation. (3)
- 4.2 A  $10 \text{ dm}^3$  steel vessel that holds a sample of oxygen gas at  $25^\circ\text{C}$  and  $100\text{kPa}$  develops a leak. Some of the oxygen gas escapes before the leak is repaired. The pressure of the  $\text{O}_2$  in the vessel after the leak is repaired is  $55\text{kPa}$ . The temperature remains at  $25^\circ\text{C}$ .

Calculate the mass of oxygen gas that leaked.

(8)  
[17]

**QUESTION 5**

A compound contains the elements carbon, hydrogen and oxygen only. It consists of 54.56% carbon and 36.36% hydrogen. The molar mass of the compound is  $132 \text{ g}\cdot\text{mol}^{-1}$ .

- 5.1 State the definition of *empirical formula*. (2)
- 5.2 Calculate the empirical formula of the compound. (6)
- 5.3 Determine the molecular formula of the compound. (2)
- [10]**

**QUESTION 6**

Iron is recovered from iron ore ( $\text{Fe}_2\text{O}_3$ ) in a blast furnace. The following reaction takes place.



In one such reaction, 160g of **impure** iron ore was reacted and  $63 \text{ dm}^3$  of  $\text{CO}_2$  was produced at STP.

- 6.1 Write down the definition of the *mole*. (2)
- 6.2 Calculate the number of  $\text{CO}_2$  molecules that formed at STP (4)
- 6.3 Calculate the maximum no of moles of iron that will be formed in the above reaction. (2)
- 6.4 Calculate the percentage purity of the iron ore sample used. (4)
- [12]**

**QUESTION 7**

Industrially, vanadium metal, (V) which is used in steel alloys can be obtained by reacting vanadium pentoxide ( $\text{V}_2\text{O}_5$ ) with calcium at high temperatures. The balanced equation for the reaction is:



During an industrial process 31850 g of  $\text{V}_2\text{O}_5$  reacts with  $2 \times 10^4$  g of Ca.

- 7.1 State the definition of a *limiting reagent*. (2)
- 7.2 Calculate the theoretical yield of vanadium. (6)
- 7.3 Calculate the percentage yield if  $8,67 \times 10^3$  g of vanadium is obtained. (2)

**[10]**



**QUESTION 8**

4.14 g of solid  $\text{LiNO}_3$  is first dissolved in a small amount of water and then made up to a certain final volume so that the concentration of the solution is  $0,05 \text{ mol.dm}^{-3}$

- 8.1 Write down the definition of *concentration*. (2)
- 8.2 Calculate the number of moles of  $\text{LiNO}_3$  used. (2)
- 8.3 Calculate the final volume of the solution. (3)
- 8.4 An additional  $250 \text{ cm}^3$  of water is now added to this solution. Calculate the new concentration of the solution (4)

**[11]****TOTAL MARKS: 100**

TABLE 3: THE PERIODIC TABLE OF ELEMENTS

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18			
(I)	(II)	(III)										(III)	(IV)	(V)	(VI)	(VII)	(VIII)			
1 H	4 Be	21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr			
3 Li	9 Be	40 Ca	41 Sc	42 Ti	43 V	44 Cr	45 Mn	46 Fe	47 Co	48 Ni	49 Cu	50 Zn	51 Ga	52 Ge	53 As	54 Se	55 Br	56 Kr		
7 Li	12 Mg	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe		
11 Na	20 Ca	39 K	40 Ca	41 Sc	42 Ti	43 V	44 Cr	45 Mn	46 Fe	47 Co	48 Ni	49 Cu	50 Zn	51 Ga	52 Ge	53 As	54 Se	55 Br	56 Kr	
19 K	20 Ca	37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe	
23 Na	24 Mg	37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe	
19 K	20 Ca	37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe	
39 K	40 Ca	37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe	
55 Cs	56 Ba	87 Fr	88 Ra	89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr		
133 Cs	137 Ba	179 Hf	181 Ta	184 W	186 Re	190 Os	192 Ir	195 Pt	197 Au	201 Hg	204 Tl	207 Pb	209 Bi	210 Po	210 At	210 Rn	210 Fr	210 Ra	210 Ac	
87 Fr	88 Ra	89 Ac	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr	103 Lr	103 Lr	103 Lr	103 Lr
58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu	71 Lu	71 Lu	71 Lu	71 Lu	71 Lu	71 Lu	71 Lu
140 Ce	141 Pr	144 Nd	144 Pm	150 Sm	152 Eu	157 Gd	159 Tb	163 Dy	165 Ho	167 Er	169 Tm	173 Yb	175 Lu	175 Lu	175 Lu	175 Lu	175 Lu	175 Lu	175 Lu	175 Lu
90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr	103 Lr	103 Lr	103 Lr	103 Lr	103 Lr	103 Lr	103 Lr
232 Th	238 U	238 U	238 Np	238 Pu	238 Am	238 Cm	238 Bk	238 Cf	238 Es	238 Fm	238 Md	238 No	238 Lr	238 Lr	238 Lr	238 Lr	238 Lr	238 Lr	238 Lr	238 Lr

TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIËSE KONSTANTES

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure <i>Standaarddruk</i>	$p^{\circ}$	$1,013 \times 10^5 \text{ Pa}$
Molar gas volume at STP <i>Molêre gasvolume by STD</i>	$V_m$	$22,4 \text{ dm}^3 \text{ mol}^{-1}$
<i>Molêre gaskonstante</i> Molar gas constant	R	$8,31 \text{ J}\cdot\text{K}^{-1}\cdot\text{mol}^{-1}$
Standard temperature <i>Standaardtemperatuur</i>	$T^{\circ}$	273 K
Avogadro's Constant	N or/ $N_A$	$6,022 \times 10^{23} \text{ mol}^{-1}$
Charge on Electron <i>Lading op elektron</i>	e	$-1,6 \times 10^{-19} \text{ C}$

TABLE 2: FORMULAE/TABEL 2: FORMULES

$n = \frac{m}{M}$	or/ of $C = \frac{n}{V}$ $C = \frac{m}{MV}$	$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$ $pV = nRT$
-------------------	---------------------------------------------------	-----------------------------------------------------------